

**Plant trees for the planet:  
the potential of forests for climate change  
mitigation and the major drivers of national  
forest area**

Sebastian Mader

Institute of Sociology  
University of Bern  
Fabrikstrasse 8  
3012 Bern  
Switzerland

Phone: +41 31 631 48 16  
Email: [sebastian.mader@soz.unibe.ch](mailto:sebastian.mader@soz.unibe.ch)

ORCID: [0000-0003-3400-4715](https://orcid.org/0000-0003-3400-4715)

03 June 2019

This is a post-peer-review, pre-copyedit version of an article published in *Mitigation and Adaptation Strategies for Global Change*. The final authenticated version is available online at: <http://dx.doi.org/10.1007/s11027-019-09875-4>.

Plant trees for the planet: the potential of forests for climate change mitigation and the major drivers of national forest area

## Abstract

Forests are one of the most cost-effective ways to sequester carbon today. Here, I estimate the world's land share under forests required to prevent dangerous climate change. For this, I combine newest longitudinal data of FLUXNET on forests' net ecosystem exchange of carbon (NEE) from 78 forest sites ( $N=607$ ) with countries' mean temperature and forest area. This straightforward approach indicates that the world's forests sequester  $8.3 \text{ GtCO}_2\text{yr}^{-1}$ . For the  $2^\circ\text{C}$  climate target the current forest land share has to be doubled to 60.0 % to sequester an additional  $7.8 \text{ GtCO}_2\text{yr}^{-1}$ , which demands less red meat consumption. This afforestation/reforestation (AR) challenge is achievable, as the estimated global biophysical potential of AR is  $8.0 \text{ GtCO}_2\text{yr}^{-1}$  safeguarding food supply for 10 billion people. Climate-responsible countries have the highest AR potential. For effective climate policies, knowledge on the major drivers of forest area is crucial. Enhancing information here, I analyse forest land share data of 98 countries from 1990 to 2015 applying causal inference ( $N=2,494$ ). The results highlight that population growth, industrialization, and increasing temperature reduce forest land share, while more protected forest and economic growth generally increase it. In all, this study confirms the potential of AR for climate change mitigation with a straightforward approach based on the direct measurement of NEE. This might provide a more valid picture given the shortcomings of indirect carbon stock-based inventories. The analysis identifies future regional hotspots for the AR potential and informs the need for fast and forceful action to prevent dangerous climate change.

**Keywords:** Forest area; climate change mitigation; carbon sequestration; net ecosystem exchange; fixed effects panel regression; FLUXNET; FAO;

**Funding sources:** This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

**Competing interests:** The author declares that he has no competing interests.

**Acknowledgements:** This work used eddy covariance data acquired and shared by the FLUXNET community, including these networks: AmeriFlux, AfriFlux, AsiaFlux, CarboAfrica, CarboEuropeIP, CarboItaly, CarboMont, ChinaFlux, Fluxnet-Canada, GreenGrass, ICOS, KoFlux, LBA, NECC, OzFlux-TERN, TCOS-Siberia, and USCCC. The ERA-Interim reanalysis data are provided by ECMWF and processed by LSCE. The FLUXNET eddy covariance data processing and harmonization was carried out by the European Fluxes Database Cluster, AmeriFlux Management Project, and Fluxdata project of FLUXNET, with the support of CDIAC and ICOS Ecosystem Thematic Center, and the OzFlux, ChinaFlux and AsiaFlux offices.

## 1 Introduction

Forests provide many tangible and intangible ecosystem services integral for human well-being (e.g. Ellison et al. 2017, Federici et al. 2015). Beyond this, forests are considered one of the most suitable ways to sequester carbon today, as afforestation and reforestation (AR) are relatively cost-effective, and associated with least expected adverse effects on biogeochemical and biogeophysical systems (Fuss et al. 2018, Griscom et al. 2017, IPCC 2014, Smith et al. 2016, Sonntag et al. 2016).

Recent global estimates on the current net carbon sink of established forests (i.e. carbon sequestration) range from 2.2 (Federici et al. 2015) to 8.0 (Grassi et al. 2018, Oleson et al. 2013)<sup>1</sup> to 8.8 gigatons of carbon dioxide per year ( $\text{GtCO}_2\text{yr}^{-1}$ ; Pan et al. 2011). Evaluations of the maximum biophysical sequestration potential of AR vary from 1.1 to 12.1  $\text{GtCO}_2\text{yr}^{-1}$  (Smith et al. 2016, Minx et al. 2018, Ciais et al. 2013). However, all these estimates are based on the calculation of changes in carbon stocks along Intergovernmental Panel on Climate Change (IPCC) guidelines (IPCC 2006) or the Houghton bookkeeping method (Houghton et al. 2012), providing an indirect and mostly incomplete measure of forests' net ecosystem exchange of carbon (NEE). This approach requires periodic information on the carbon content of biomass, and involves fundamental assumptions on carbon stocks – especially when reliable data is missing. This is notably true for many developing nations (Grassi et al. 2018, IPCC 2006). Moreover, each of these country estimates is based on different data quality, definitions of forest area, and accounting methods. Though data quality is gradually improving, this suggests a sizable challenge to develop a valid and internationally comparable inventory of global forest carbon fluxes based on indirect stock-based techniques (Grassi et al. 2018).

---

<sup>1</sup> Results from the simulations of the Dynamic Global Vegetation Model (DGVM) Community Land Model (CLM) version 4.5 (Oleson et al. 2013; Table SI 8 in Grassi et al. 2018);

This study has four objectives: First, I provide estimates of the annual carbon sequestration of established forests, and the biophysical climate change mitigation potential of AR based on the direct micrometeorological measurement of NEE as provided by FLUXNET (NASA 2015) (section 2). With this direct measurement of above canopy carbon flux no information on carbon stocks is needed to infer NEE. Thus, NEE estimates based on FLUXNET data may provide a more valid picture of forests' carbon sink and their mitigation potential. Second, with this straightforward approach, I infer the forest land share required to meet the 2 °C climate target and three AR scenarios to acquire this goal (section 3; see Appendix A Methods and Materials for details). Third and subsequently, I identify the countries with the largest climate liabilities, and economic capabilities while having the greatest mitigation potential through AR (section 4).

Fourth, for effective policies targeted at enhancing forests and climate change mitigation, knowledge on the key drivers of forest area is essential. However, information on causal relationships of forest gain and loss is sparse, and unconsolidated (Aguilar and Song 2018, Morales-Hidalgo et al. 2015) with a focus on forest loss (Busch and Ferretti-Gallon 2017). Yet, this is only half of the story to be told. Thus, here I identify the major predictors of the forest land share of 98 countries from 1990 to 2015 gathered from the Food and Agriculture Organization of the United Nations (FAO 2018) applying causal inference (section 5). The last section summarizes and discusses the main results, and closes with some concluding remarks.

## 2 Global and regional forest carbon sink

To quantify the NEE of countries' forests, I utilize the newest available micrometeorological FLUXNET data of 78 measurement towers in forests of 16 countries on five continents from 2000 to 2014 ( $N=607$ ; Table B.1 in Appendix B Supplementary Figures and Tables). Multiple linear ordinary least squares (OLS) regression identifies annual mean temperature as the main determinant of forests' NEE (u-shaped relationship) in this data (Table B.2, and Figure B.1 in Appendix B). Model predictions on countries' NEE of forests using countries' average temperature taken from the World Bank (2018) show that established forests sequester  $-8.8 \text{ tCO}_2\text{ha}^{-1}\text{yr}^{-1}$  on average in 2015 (median:  $-9.2$ ; Appendix A). This is rather close to prior assessments based on indirect measurements of NEE (Sohngen 2010). Portugal has the highest negative NEE with a net absorption of  $-15.1 \text{ tCO}_2\text{ha}^{-1}\text{yr}^{-1}$ , whereas the highest positive NEE is observed for Canada with a net release of  $16.3 \text{ tCO}_2\text{ha}^{-1}\text{yr}^{-1}$  (Figure 1a). The forests of almost all countries are net absorbers of carbon, except the boreal forests in Canada, the Russian Federation, and Mongolia that are net sources of carbon. This might be due to diebacks of these boreal forests resulting from insect outbreaks and wildfires due to higher mean temperatures and droughts induced by climate change (Canadell and Raupach 2008). As introspection of Figure 1a reveals, NEE varies by climate forest domain following a u-shaped mean temperature – NEE relationship. The carbon sequestration of boreal forests is lowest with a mean NEE of  $-1.1 \text{ tCO}_2\text{ha}^{-1}\text{yr}^{-1}$ , while it is highest for temperate forests with  $-12.6 \text{ tCO}_2\text{ha}^{-1}\text{yr}^{-1}$ . Tropical forests' NEE lies in-between with an average of  $-6.0 \text{ tCO}_2\text{ha}^{-1}\text{yr}^{-1}$ . This pattern is in line with former research (Brumme et al. 2005).


Multiplying countries' average NEE per hectare by their forest area (FAO 2018, Figure 1b) suggests an overall forest carbon sink of  $-8.3 \text{ GtCO}_2\text{yr}^{-1}$  or  $-1.1 \text{ tCO}_2\text{yr}^{-1}$  per

capita (p.c., UNPD 2017) in 2015. Carbon sequestration is highest in the forests of the United States and Brazil ( $-3.2 \text{ GtCO}_2\text{yr}^{-1}$  each), followed by China ( $-2.0 \text{ GtCO}_2\text{yr}^{-1}$ ), Australia ( $-1.5 \text{ GtCO}_2\text{yr}^{-1}$ ) and the Democratic Republic of the Congo ( $-1.1 \text{ GtCO}_2\text{yr}^{-1}$ ). The rest of the world's countries has a net absorption of less than  $-1.0 \text{ GtCO}_2\text{yr}^{-1}$  each, and Canada, Mongolia, and the Russian Confederates have a substantial net release of  $16.7 \text{ GtCO}_2\text{yr}^{-1}$  in sum.

The global estimate of this rather simple approach using direct carbon flux measurements of NEE is fairly close to the estimates of two recent studies applying more complicated, indirect, carbon stock-based inventories of NEE (Grassi et al. 2018, Oleson et al. 2013, Pan et al. 2011). Grassi et al. (2018) report a global forest carbon sink of  $-8.0 \text{ GtCO}_2\text{yr}^{-1}$  for the Community Land Model (version 4.5; Oleson et al. 2013)<sup>2</sup> and Pan et al. (2011) estimate a sink of  $-8.8 \text{ GtCO}_2\text{yr}^{-1}$  based on changes in carbon stocks.

---


<sup>2</sup> This Dynamic Global Vegetation Model (DGVM) could be considered one of the most elaborate DGVMs as it comprises the most relevant ecological characteristics as compared to other commonly used DGVMs (Table SI 7 in Grassi et al. 2018).



**Figure 1 | Net ecosystem exchange (NEE) of CO<sub>2</sub> of countries' forests in 2015. a-c**, Data source for the calculation of NEE of CO<sub>2</sub> of countries' forests is FLUXNET (NASA 2015), World Bank (2018) and FAO (2018). Negative numbers indicate net absorption of carbon, positive numbers its net release. **a**, Carbon sequestration in tCO<sub>2</sub>ha<sup>-1</sup>yr<sup>-1</sup> of forest area (mean = -8.8, median = -9.2, min. (Portugal) = -15.1, max. (Canada) = 16.3). **b**, Countries' overall forest carbon sequestration in GtCO<sub>2</sub>yr<sup>-1</sup> (sum = -8.3 GtCO<sub>2</sub>yr<sup>-1</sup>). **c**, Countries' overall NEE potential of afforestation/reforestation (AR) in GtCO<sub>2</sub>yr<sup>-1</sup> based on scenario 3 exceeding the 7.8 GtCO<sub>2</sub>yr<sup>-1</sup> required to meet the 2 °C respectively 3 tCO<sub>2</sub> per capita climate target (sum = -8.0 GtCO<sub>2</sub>yr<sup>-1</sup>; see text and Appendix A for details).

### 3 Forest land share trends and AR scenarios

Before evaluating countries' climate change mitigation potential of AR (Figure 1c), the current forest land share, suitable land for AR as well as competing land uses have to be quantified. The average forest land share as provided by the FAO shrunk from 31.8 % in 1990 to 30.8 % in 2015 (Figure 2), which corresponds to a forest loss of 1.3 Mkm<sup>2</sup> – an area as large as Peru.



**Figure 2 | Forest land share in international comparison 1990 and 2015.** Depicted are the top and bottom five countries, top five countries with respect to overall forest area (FAO 2018) by climate domain in 2015, and members of the G7 and BRIICS if not already included. Dark blue solid line = mean 2015; Grey solid line = mean 1990; Scenario 1: Red solid line = required forest share for the 2 °C climate target, red long-dashed line = achievable forest share; Scenario 2: red dashed line = achievable forest share; Scenario 3: Green solid line = required forest share, green short-dashed line = achievable forest share. See text and Appendix A for details.

As Figure 2 shows, European countries like France, Italy, Germany, and Norway resemble the mean of 2015. The forest land share varies strongly: Laos ranks highest with 81.3 % and is followed by Papua New Guinea, Finland, Guinea-Bissau, and Sweden constituting the top five. The bottom five countries with almost no forests are Algeria, Saudi Arabia, Mauritania, Libya, and Egypt. Between 1990 and 2015 Indonesia incurred the greatest loss of almost a quarter and Brazil as top carbon sequestering country lost 10.0 % of its tropical forests. The greatest gain was



accomplished by China with a one third increase in forest area while ranking third in overall NEE. For the United States as top carbon absorbing nation, almost no change in forest cover was observed in this period.

Furthermore, Figure 2 presents the required as well as the achievable forest land share of three different AR scenarios to prevent dangerous climate change. The global annual gross carbon budget to fulfil the 2 °C climate target with a probability of at least 66 % is an estimated 30 GtCO<sub>2</sub> (IPCC 2014, Friedlingstein et al. 2014, Meinshausen et al. 2009). Assuming an average annual world population of 9.8 billion people until 2100 (UNPD 2017), this goal translates into ~3 t of gross CO<sub>2</sub> emissions per capita (p.c.) and year.

First, scenario 1 is the baseline scenario. It assumes constant production and consumption patterns, constant other carbon sinks, and a further required emissions reduction of 1.0 tCO<sub>2</sub>yr<sup>-1</sup> p.c. after accounting for the overall forest carbon sequestration of 0.8 tCO<sub>2</sub>yr<sup>-1</sup> p.c.. Hence, in scenario 1 the required forest land share to meet the 2 °C respectively the 3 tCO<sub>2</sub> p.c. climate target is 67.8 % (red solid line in Figure 2) to additionally sequester 9.8 GtCO<sub>2</sub>yr<sup>-1</sup> (Appendix A). The red long-dashed line is the forest land share that can be achieved via 100 % AR of all shrub-covered areas and herbaceous vegetation as retrieved from the FAO (2018; 44.8 % forest land share). This is more than one third of the required AR. Second, in scenario 2 a forest land share of up to 57.5 % can be achieved by additionally afforesting and reforesting 44 % of permanent grassland and cropland (FAO 2018), assuming current diets and an average land demand of 2,100 m<sup>2</sup> p.c. (Hallström et al. 2015) for feeding an expected 9.8 billion people per year (red dashed line in Figure 2). This represents more than two thirds of this tremendous AR challenge. Finally, in scenario 3 healthier diets with reduced red and ruminant meat consumption decrease agricultural land demand further by 28.0 % to 1,510 m<sup>2</sup> p.c. and reduce dietary-related emissions by 0.2 tCO<sub>2</sub>yr

<sup>1</sup> p.c. (Hallström et al. 2015). This yields a required forest land share of 60.0 % to meet the 2°C climate target (green solid line in Figure 2) equivalent to an additional 7.8 GtCO<sub>2</sub>yr<sup>-1</sup> to be sequestered by forests. Thus, in this healthy diet scenario further AR of grassland and cropland results in an attainable 62.0 % of forest land share (8.0 GtCO<sub>2</sub>yr<sup>-1</sup>; green short-dashed line in Figure 2).

Consequently, the 2 °C climate target can be met by almost doubling the current forest area whilst safeguarding food security with a healthy diet. This outstanding challenge means 37.9 Mkm<sup>2</sup> more of forest area or an estimated 2.6 trillion additional trees. Approximately, this corresponds to the number of trees lost since the start of human civilization (Crowther et al. 2015). This challenge translates into approximately 260 trees p.c. or one tree p.c. per week for a realisation time of five years.


Realizing the need for large-scale AR, there are promising worldwide projects like 'Plant for the Planet', which aims at planting one trillion trees. Since 2007, this project has planted 13.6 billion trees (Plant for the Planet 2019) – 0.5 % of the climate target. In 2017 the World Wildlife Fund, the Wildlife Conservation Society and BirdLife International launched the 'Trillion Trees' program aiming at restoring one trillion trees by 2050 (Trillion Trees 2019). Furthermore, the 'Bonn Challenge' strives for the restoration of 3.5 Mkm<sup>2</sup> of forests by 2030 (~9.2 % of AR required for the 2 °C target). To date pledges exceed 1.7 Mkm<sup>2</sup> (International Union for Conservation of Nature 2019). To achieve the targets of all three voluntary initiatives together would account for the vast majority of the required AR (86 %). 260 trees per capita seems a relatively low number. However, the need for fast and forceful AR is high leaving this venture an ambitious challenge.

#### 4 Liabilities, AR potentials, and capabilities

Given that call, who is in charge of action? Being the country with the highest negative NEE of established forests (Figure 1b), and the world's second largest carbon emitter (Janssens-Maenhout et al. 2017), the United States of America rank highest in the climate change mitigation potential of countries through AR (NEE =  $-1.0 \text{ GtCO}_2\text{yr}^{-1}$ ; Figure 1c). Figure 1c also demonstrates that the world's largest carbon emitter and third largest carbon absorber in forests, China, has the second highest AR potential ( $-0.8 \text{ GtCO}_2\text{yr}^{-1}$ ). This offers a great opportunity for the United States, and China, accounting for almost half of the global carbon emissions and having to bear one of the highest domestic social costs of carbon emissions (Ricke et al. 2018), to take their responsibility for climate change mitigation seriously. Together with Australia, Argentina, and Brazil they form the top five countries with respect to mitigation potential through AR, accounting for almost half of its total.

The radar plots in Figure 3 provide a more comprehensive picture of the countries' climate change liabilities, forests' mitigation contributions, AR potentials, and economic capabilities for action in worldwide comparison. One group of countries at the top of the ranking of the sum score of these characteristics is formed by those ranking highest in mitigation potential of AR, while being among the largest emitters of CO<sub>2</sub> (p.c.) and the wealthiest nations (Figure 3a-c,e-j,n). These countries are Japan, Spain, France, Australia, the United States, Argentina, Italy, Germany, Brazil, and the United Kingdom. Hence, these states could take over their responsibility for climate change mitigation relatively easily via large-scale domestic AR activities. Figure 2 indicates that the forest land share of three of these countries, France, Italy, and the United Kingdom, grew between 1990 and 2015, while Brazil, and Argentina experienced forest loss.

Another group of nations is both liable of global warming and has high AR potential, but to some extent lacks economic strength to implement large-scale measures. Countries like China, Peru, South Africa, Indonesia, and India fall into this group (Figure 3k-m,o,p). Indonesia and Peru reflect this, since these countries lost forests between 1990 and 2015 (Figure 2). By contrast, China, and India gained forest in this period probably due to large-scale AR programs. These nations and poor countries with little climate responsibility but large AR potential like the Democratic Republic of the Congo (Figure 3t) need multilateral financial assistance, foremost from wealthy, climate-responsible states, to unfold their AR potential. This applies to countries, which additionally have relatively low or no AR mitigation potential like South Korea, Sweden, Canada, and the Russian Federation (Figure 3d, q-s). This could be a worthwhile enhancement of the REDD+ (Reducing Emissions from Deforestation and Forest Degradation) framework.



**Figure 3 | Country ranking of climate responsibility, forests' mitigation contribution and potential, and economic capabilities in 2015.** a-y, Radar plots of countries' relative performance with regard to climate responsibility (CO<sub>2</sub>, and CO<sub>2</sub> per capita (p.c.) emissions (Janssens-Maenhout et al. 2017)), forests' mitigation contribution (forest land share (%; FAO 2018), net ecosystem exchange (NEE) per ha, and national NEE), forests' mitigation potential (NEE potential)), and economic capabilities (gross domestic product (GDP) p.c. (IMF 2018)). The numbers 1 to 5 on the spokes of the radars indicate the quintile the country ranks (1 = lowest, 5 = highest). The numbers in the centre of each radar represent the sum of quintiles of each country. Presented are the top and bottom five countries with respect to this sum, the top five countries of overall forest area by climatic forest domain and members of the G7 and BRIICS. The full country ranking of the sum score and all included variables can be obtained from Table B.4 in Appendix B.

## 5 Predictors of national forest land share

Nonetheless, the plea for international cooperation and referring to climate change responsibility is not enough. For effective policies targeted at the enhancement of forests, profound knowledge on the key drivers of national forest area is crucial. Previous research has focused on determinants of forest loss with different regional and temporal cover and a focus on satellite-derived data in recent years (Busch and Ferretti-Gallon 2017, Leblois et al. 2017). However, these studies are agnostic about AR and forest regrowth, as some authors critically remark themselves (DeFries et al. 2010). Focusing on forest loss only shines light on half of the story to be told. Hence, causal information on the predictors of national forest land share analysing panel data of many countries by means of causal inference is still sparse and unconsolidated (Aguilar and Song 2018, Morales-Hidalgo et al. 2015).


Aguilar and Song (2018), and Morales-Hidalgo et al. (2015) are the only two studies regressing changes in national forest area as provided by the FAO on changes in countries' socioeconomic characteristics utilizing fixed effects (FE) panel regression models. Morales-Hidalgo et al. (2015) is the first study regressing national forest area between 1990 and 2015 gathered from the FAO on a few socio-economic and political indicators applying causal inference. The results of their country and year FE panel regression models (Table 6 in Morales-Hidalgo et al. 2015) suggest that population growth reduces forest area, whereas GDP p.c. and protected areas increase it. Nonetheless, the results of Morales-Hidalgo et al. (2015) could be biased by omitting other substantial drivers of forest land share. Aguilar and Song (2018) is the only study analysing the ratio between national forest area and land area (i.e. forest land share) ensuring comparability of changes in forest cover between countries irrespective of their total land area. In their FE models, Aguilar and Song (2018) include agricultural land area, 10-year lagged GDP growth rate, GNI p.c., population growth rate,

population density, share of rural population, rate of secondary school enrolment, its 15-year lagged values, and the squares of all these characteristics as independent variables. The results of their beta-logistic generalized linear mixed models with ratio response indicate that all of the considered covariates are substantially related to forest land share (Table 3 in Aguilar and Song 2018). However, FE models including both levels and lags of the same characteristics produce biased results, if the causal effects emerge immediately (Vaisey and Miles 2017), as it is the case in Aguilar and Song (2018). Furthermore, the results of Aguilar and Song (2018) could be biased by omitting important confounding variables.

To improve, consolidate and expand previous studies, here I regress the forest land share of 98 countries from 1990 to 2015 as provided by the FAO on socio-economic, political, and ecological characteristics applying country and year FE regression models (Brüderl and Ludwig 2015; Appendix A). The 98 countries analysed (Table B.7) have high or sufficient quality of forest area data (tiers 3 and 2; FAO 2016) and comprise around 89 % of global forest area in 2015 (Keenan et al. 2015). All other countries, which have unreliable data solely based on expert estimates (tier 1) are excluded from the analysis.

First, one of the best-documented drivers of deforestation is agricultural expansion (Jorgenson 2006). As model 1 of Figure 4 shows, a 1 % within-country increase in agricultural land share on average leads to a 0.2 % within-country decrease in the forest land share. Population growth explains this effect, as it disappears when population size is included in the regression (model 2). Population growth of 1 % yields deforestation of 0.27 %. This suggests that agricultural expansion allows population growth, which in turn exerts pressure on forests because of land demands for housing,

mobility, and other resources.<sup>3</sup>



**Figure 4 | Predictors of national forest land share.** Coefficient plots of unstandardized regression coefficients (dark blue filled circles) of country and year fixed effects regressions of national forest land share on various successively included predictors (models 1-6) including 95 % confidence intervals (dark blue bars; see Table B.5 in Appendix B for details). All six models contain dummy variables for each year to control for overall time-trends. All variables are included by taking their natural logarithm allowing the estimation of elasticities. ‘n’ refers to the number of countries, and ‘N’ to the number of observations (number of countries (n) times the number of years). Table B.6 in Appendix B describes all variables and Table B.7 lists all countries included in the models.

Second, it has often been hypothesized that urbanization slows deforestation and promotes AR, because the per capita land demand of cities is assumed to be lower as compared to rural areas (Jorgenson 2006). However, the models 2-6 of Figure 4 reveal that increasing rates of the population living in urban areas are not substantially related to countries’ forest area.

Third, the direction of the impact of growing wealth on forest cover is widely discussed in the literature (e.g. Jorgenson 2006). There has been widespread consent that deforestation activities prevail at low and middle levels of gross domestic product (GDP) p.c. and AR activities outweigh deforestation at higher levels of GDP p.c. following a trajectory referred to as the environmental Kuznets curve (EKC; Aguilar and

<sup>3</sup> Moreover, in the FE regression of population ( $N=2504$ ,  $n=98$ ), the elasticity of agricultural land is 0.49 ( $p < 0.001$ ). Together with the results of the models 1 and 2 of Figure 4, this suggests that population growth mediates the relationship between agricultural expansion and forest loss.



Song 2018). However, the empirical evidence for a forest EKC is mixed and the two most recent and elaborate studies found evidence for a clear positive relationship between GDP and forest cover invalidating the forest EKC hypothesis (Aguilar and Song 2018, Morales-Hidalgo et al. 2015). Models 3-6 of Figure 4 highlight this as well: Economic growth of 1 % increases forest land share by 0.1 % irrespective of economic structure and wealth levels.<sup>4</sup> Hence, this supports the notion that wealth at least to some extent leads to more awareness for the ecosystem services of forests and the need to protect them.

In addition to that and extending prior studies, economic structural change could affect forest transition net of GDP growth. An increasing GDP share of the industry sector might introduce pressure on forestlands because of relatively high land requirements of industrial production sites and higher returns for industrial production than for forest products. As models 3-6 of Figure 4 indicate, there is some evidence in favour of this argument, because a 1 % increase in the GDP share of the industry sector yields a 0.1 % decrease in forest land share. In turn, expansion of the service sector could release pressure from forests, as services are presumed to have less land demand. However, in the data there is no support for this notion, since a 1 % increase in the GDP share of the service sector is also related to a 0.1 % decline in forest cover. Yet, this effect is not statistically significant at the  $p = 0.05$  level.

Furthermore, the model is enhanced by including an indicator of foreign trade in forest products. It has been a common concern that forest products trade could be one of the reasons for deforestation especially in poor countries with tropical forests and few alternatives of employment to timber logging or farming. By contrast, one can argue that foreign trade of forest products could be an incentive for forest conservation,

---

<sup>4</sup> The partial residual plot for GDP of a penalized splines FE regression (Ruppert et al. 2003) adequately modelling non-linearities confirms this, too (Figure B.2 in Appendix B).

when the net return of forestry investments and sustainable forest management is greater than the net return for forest clearing for agricultural production (Burgess 1993). However, models 4-6 of Figure 4 demonstrate that increases in exports of forest products relative to their imports do not substantially alter countries' forest area.

Moreover, policies for forest protection may contribute to stop deforestation and forest degradation, and foster AR activities with the aim of enhancing the global forest carbon sink, conserving biodiversity, and safeguarding other ecosystem services of forests. These goals are part of manifold international initiatives and agreements on forest protection. Designating and managing protected areas has been a primary strategy to achieve these goals (Morales-Hidalgo et al. 2015). Hence, protected forest area serves as an indicator for a country's willingness to sustain the ecosystem services of forests and to commit to AR activities. As models 5 and 6 of Figure 4 reveal, a 1 % increase in protected forest area is associated with forest growth of 0.06 %. This effect is statistically significant, but rather small. This is in line with the results of Morales-Hidalgo et al. (2015).

Finally, climate change itself might harm forest ecosystems leading to forest degradation and forest loss. Long-term case studies of tree mortality indicate that higher mean temperature and droughts increase tree mortality and the frequency of wildfires (Canadell and Raupach 2008, Young et al. 2017, Martin 2015). However, it is still unclear whether this also applies to forest loss on a global scale. As model 6 of Figure 4 shows, a 1 % increase in countries' mean air temperature reduces their forest area on average by 0.1 %, while severe drought events do not affect forest cover immediately and *ceteris paribus*. This suggests that global warming contributes to forest loss, even though the effect is rather small.

## 6 Discussion and Conclusion

Altogether, this study suggests that dangerous climate change could be prevented solely by AR, as forests' biophysical climate change mitigation potential safeguarding food security with healthy diets (scenario 3) exceeds the required additional carbon uptake for the 2 °C target. For this, the study estimates countries' carbon sequestration of forests based on the direct micrometeorological measurement of NEE, average temperature and forest area. This straightforward, direct carbon flux-based method provides estimates that are comparable to the most recent studies applying more complicated, indirect carbon stock-based inventories of NEE. The direct approach followed here might provide a more valid picture given the outlined shortcomings of indirect carbon stock-based inventories. However, the direct approach rests on the assumption that countries' average temperature is a valuable approximation of the mean climatic conditions of their forests. Moreover, uncertainties stem from data gaps on the NEE of tropical forest biomes, as Figure B.1 in Appendix B demonstrates. Further uncertainties may arise from varying tree density, age, species, species richness and the health of forests (Hawes 2018). Hence, further validation of these initial findings is needed. This includes the establishment of additional and more precise FLUXNET measurement towers especially in tropical forests to close data gaps, and to increase accuracy and spatial resolution of model predictions.

Furthermore, the analysis identifies future regional hotspots for the AR potential. The United States, China, Australia, Argentina, and Brazil are the top five countries with respect to mitigation potential through AR, accounting for almost half of its total. However, to unfold the AR potential effectively, it is vital to establish a global mandatory carbon certificate market incorporating the forest carbon sink of countries and private forest owners. This generates financial incentives to restore and sustain forest biomes

(Sohngen 2010). Enriching voluntary initiatives like REDD+ with countries' AR potentials, climate-liabilities, and economic capabilities might be a valuable starting point for that.

Evenly important, the analysis of the major drivers of countries' forest land share highlights that curbing agricultural expansion and population growth may be a focus for AR policies. Moreover, forests' vulnerability to global warming points to the necessity to plant the right trees in the right places. Therefore, sustainable regional forest management needs to identify the tree species most resilient to temperature increases, and enhance the biodiversity of forests (Huang et al. 2018, Liang et al. 2017). Together with growing wealth, the expansion of protected forest areas is a suitable way to amplify the forest carbon sink, conserve biodiversity, and safeguard other vital ecosystem services provided by forests.

Nevertheless, biophysical, social, and economic challenges alongside large-scale AR might jeopardize its potential benefits (e.g. Canadell and Raupach 2008, Smith et al. 2016, Fuss et al. 2018), and contest the feasibility of the three presented AR scenarios. In general, all three presented AR scenarios a priori exclude land cover types that are, by themselves, biophysically unsuitable for near-term and cost-efficient AR (i. e. artificial surfaces, permanent snow and glaciers, terrestrial barren land, and sparsely natural vegetated areas). In addition, all scenarios safeguard food supply for 10 billion people. However, the feasibility of all three scenarios more or less depends on the socio-economic pressure exerted on the land designated to be afforested/reforested. Griscom et al. (2017) report that almost half of the existing AR potential could be cost-effectively realized below US\$100 tCO<sub>2</sub><sup>-1</sup> (the estimated social cost of 1 tCO<sub>2</sub> emitted within the 2 °C climate target). More than 10 % of the AR potential are achievable at low cost (<US\$10 tCO<sub>2</sub><sup>-1</sup>). At least part of scenario 1, the AR of shrub-covered and herbaceous vegetation, might be reachable at low cost.

However, costs are expected to be higher for the AR of agricultural land (permanent grassland and cropland; scenarios 2 and 3). Agricultural expansion and increases in population density increase the opportunity costs of not clearing forests and the costs of AR, and decrease forest cover (as shown in this study). Near-term costs might be even higher, when a large-scale diet transition away from red and ruminant meat is demanded to free up additional land for AR (scenario 3). Yet, meat reduced diets are regarded as 'win-win diets' fostering both public health and the environment in the long-run (Willett et al. 2019). Moreover, and as this study demonstrates, the growing wealth of nations decreases the relative costs of AR and conveys forest protection and AR. Nonetheless, well-tailored AR policies have to account for possible trade-offs between climate change mitigation through AR and benefits for the local population. Here, agroforestry and policies targeted at the promotion of timber as building material whilst substituting carbon-intensive concrete and steel could be especially beneficial, and may substantially promote climate change mitigation (Oliver et al. 2014, Tollefson 2017).

All told, permanent carbon storage is a prerequisite to outpace the burning of fossil carbon and reduce the CO<sub>2</sub> concentration in the atmosphere. Hence, it is vital to combine sustainably managed, large-scale AR activities with technologies for permanent carbon storage like bioenergy with carbon capture and storage (BECCS) at the end of the trees' life cycle for effective climate change mitigation (Fuss et al. 2018, Smith et al. 2016). What is more, abating emissions and applying other negative emissions technologies are valuable in order to hedge the impact of potential side effects of one mitigation option like AR (Minx et al. 2018, Sohngen 2010, Fuss 2010) to keep up with the need for fast and forceful action to prevent dangerous climate change.

## References

- Aguilar FX, Song N (2018) Forest cover, agricultural, and socio-economic development: A weighted beta-logistic approach with ratio response. *Forest Sci* 64: 129-138.
- Brüderl J, Ludwig V (2015) Fixed-effects panel regression. In: Best H, Wolf C (eds) *The SAGE handbook of regression analysis and causal inference*. SAGE, London, pp 327-358.
- Brumme R, Verchot LV, Martikainen PJ, Potter CS (2005) Contribution of trace gases nitrous oxide (N<sub>2</sub>O) and methane (CH<sub>4</sub>) to the atmospheric warming balance of forest biomes. In: Griffiths H, Jarvis PG (eds) *The Carbon Balance of Forest Biomes*. Taylor & Francis, Scarborough, Canada, pp 313-341.
- Burgess JC (1993) Timber production, timber trade and tropical deforestation. *Ambio* 22: 136-143.
- Busch J, Ferretti-Gallon K (2017) What drives deforestation and what stops it? A meta-analysis. *Rev Env Econ Policy* 11: 3-23.
- Canadell G, Raupach MR (2008) Managing forests for climate change mitigation. *Science* 320: 1456-1457.
- Ciais P, Sabine C, Bala G, Bopp L, Brovkin V, Canadell J, Chhabra A, DeFries R, Galloway J, Heimann M, Jones C, Le Quéré C, Myneni RB, Piao S, Thornton P (2013) Carbon and other biogeochemical cycles. In: Stocker TF, Qin D, Plattner G-K, Tignor MMB, Allen SK, Boschung J, Nauels A, Xia Y, Bex V, Midgley PM (eds) *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge, UK, pp 465-570.
- Crowther TW, Glick HB, Covey KR, Bettigole C, Maynard DS, et al. (2015) Mapping tree density at a global scale. *Nature* 525: 201-205.
- DeFries R, Rudel T, Uriarte M, Hansen M (2010) Deforestation driven by urban population growth and agricultural trade in the twenty-first century. *Nat Geosci* 3: 178-181.
- Ellison D, Morris CE, Locatelli B, Sheil D, Cohen J, et al. (2017) Trees, forests and water: Cool insights for a hot world. *Global Environ Chang* 43: 51-61.
- FAO – Food and Agriculture Organization of the United Nations (2016) *Global Forest Resources Assessment 2015. How are the world's forests changing?* 2<sup>nd</sup> edn. <http://www.fao.org/3/a-i4793e.pdf>. Accessed 28 May 2019.
- FAO – Food and Agriculture Organization of the United Nations (2018) FAOSTAT. <http://www.fao.org/faostat/en/#data>. Accessed 28 May 2019.
- Federici S, Tubiello FN, Salvatore M, Jacobs H, Schmidhuber J (2015) New estimates of CO<sub>2</sub> forest emissions and removals: 1990-2015. *Forest Ecol Manag* 352: 89-98.
- Friedlingstein P, Andrew RM, Rogelj J, Peters GP, Canadell JG, Knutti R, Luderer G, Raupach MR, Schaeffer M, van Vuuren DP, Le Quéré C (2014) Persistent growth of CO<sub>2</sub> emissions and implications for reaching climate targets. *Nat Geosci* 7: 709-715.
- Fuss S (2010) Forestry Carbon Sequestration. In: Lomborg B (ed) *Smart solutions to climate change*. Cambridge University Press, Cambridge, UK, pp 133-141.
- Fuss S, Lamb WF, Callaghan MW, Hilaire J, Creutzig F, et al. (2018) Negative emissions – Part 2: Costs, potentials and side effects. *Environ Res Lett* 13: 063002.
- Grassi G, House J, Kurz WA, Cescatti A, Houghton RA, et al. (2018) Reconciling global-model estimates and country reporting of anthropogenic forest CO<sub>2</sub> sinks. *Nat Clim Change* 8: 914-920.
- Griscom BW, Adams J, Ellis PW, Houghton RA, Lomax G, et al. (2017) Natural climate solutions. *P Natl Acad Sci USA* 114: 11645-11650.

- Hallström E, Carlsson-Kanyama A, Börjesson P (2015) Environmental impact of dietary change: A systematic review. *J Clean Prod* 91: 1-11.
- Hawes M (2018) Planting carbon storage. *Nat Clim Change* 8: 556-558.
- Houghton RA, House JI, Pongratz J, van der Werf GR, DeFries RS, Hansen MC, Le Quéré C, Ramankutty N (2012) Carbon emissions from land use and land-cover change. *Biogeosciences* 9: 5125-5142.
- Huang Y, Chen Y, Castro-Izaguirre N, Baruffol M, Brezzi M, et al. (2018) Impacts of species richness on productivity in a large-scale subtropical forest experiment. *Science* 362: 80-83.
- IMF – International Monetary Fund. World Economic Outlook Databases (2018). <http://www.imf.org/external/ns/cs.aspx?id=28>. Accessed 28 May 2019.
- International Union for Conservation of Nature (2019) The Bonn Challenge. <http://www.bonnchallenge.org/>. Accessed 28 May 2019.
- IPCC – Intergovernmental Panel on Climate Change (2006) 2006 IPCC guidelines for national greenhouse gas inventories. Volume 4. Agriculture, forestry and other land use. Eggleston S, Buendia L, Miwa K, Ngara T, Tanabe K (eds) Institute for Global Environmental Strategies, Hayama, Japan.
- IPCC – Intergovernmental Panel on Climate Change (2014) Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Core Writing Team, Pachauri RK, Meyer LA (eds). IPCC, Geneva.
- Janssens-Maenhout G, Crippa M, Guizzardi D, Muntean M, Schaaf E, Olivier JGJ, Peters JAHW, Schure KM (2017) Fossil CO<sub>2</sub> and GHG emissions of all world countries. Publications Office of the European Union, Luxembourg.
- Jorgenson AK (2006) Unequal ecological exchange and environmental degradation: A theoretical proposition and cross-national study of deforestation, 1990-2000. *Rural Sociol* 71: 685-712.
- Keenan RJ, Reams GA, Achard F, de Freitas JV, Grainger A, Lindquist E (2015) Dynamics of global forest area: Results from the FAO Global Forest Resources Assessment 2015. *Forest Ecol Manag* 352: 9-20.
- Leblois A, Damette O, Wolfersberger J (2017) What has driven deforestation in developing countries since the 2000s? Evidence from new remote-sensing data. *World Dev* 92: 82-102.
- Liang J, Crowther TW, Picard N, Wiser S, Zhou M, et al. (2017) Positive biodiversity-productivity relationship predominant in global forests. *Science* 354: 196.
- Martin C (2015) On the edge. The state and fate of the world's tropical rainforests. Greystone, Vancouver.
- Meinshausen M, Meinshausen N, Hare W, Raper SCB, Frieler K, Knutti R, Frame DJ, Allen MR (2009) Greenhouse-gas emission targets for limiting global warming to 2 °C. *Nature* 458: 1158-1162.
- Minx JC, Lamb WF, Callaghan MW, Fuss S, Hilaire J, et al. (2018) Negative emissions - part 1: Research landscape and synthesis. *Environ Res Lett* 13: 063001.
- Morales-Hidalgo D, Oswalt SN, Somanathan E (2015) Status and trends in global primary forest, protected areas, and areas designated for conservation of biodiversity from the Global Forest Resources Assessment 2015. *Forest Ecol Manag* 352: 68-77.
- NASA – National Aeronautics and Space Administration of the United States (2015) FLUXNET 2015 dataset (2015). <http://fluxnet.fluxdata.org/data/fluxnet2015-dataset/>. Accessed 28 May 2019.
- Oleson KW, Lawrence DM, Bonan GB, Drewniak B, Huang M, et al. (2013) Technical description of version 4.5 of the Community Land Model (CLM). <http://dx.doi.org/10.5065/D6RR1W7M>.
- Oliver CD, Nassar NT, Lippke BR, McCarter JB (2014) Carbon, fossil fuel, and biodiversity mitigation with wood and forests. *J Sustain Forest* 33: 248-275.

- Pan Y, Birdsey RA, Fang J, Houghton R, Kauppi PE, et al. (2011) A large and persistent carbon sink in the world's forests. *Science* 333: 988-993.
- Plant for the Planet (2019) Plant for the planet. [www.plant-for-the-planet.org/](http://www.plant-for-the-planet.org/). Accessed 28 May 2019.
- Ricke K, Drouet L, Caldeira K, Tavoni M (2018) Country-level social cost of carbon. *Nat Clim Change* 8: 895-900.
- Smith P, Davis SJ, Creutzig F, Fuss S, Minx J, et al. (2016) Biophysical and economic limits to negative CO<sub>2</sub> emissions. *Nat Clim Change* 6: 42-50.
- Sohngen B (2010) Forestry carbon sequestration. In: Lomborg B (ed) *Smart solutions to climate change*. Cambridge University Press, Cambridge, UK, pp 114-132.
- Sonntag S, Pongratz J, Reick CH, Schmidt H (2016) Reforestation in a high-CO<sub>2</sub> world - Higher mitigation potential than expected, lower adaptation potential than hoped for. *Geophys Res Lett* 43: 6546-6553.
- Tollefson J (2017) The wooden skyscrapers that could help to cool the planet. *Nature* 545: 280-282.
- Trillion Trees (2019) Trillion Trees. <https://www.trilliontrees.org/>. Accessed 28 May 2019.
- UNPD – United Nations Population Division (2017) *World Population Prospects 2017*. <https://esa.un.org/unpd/wpp/>. Accessed 28 May 2019.
- Vaisey S, Miles A (2017) What you can – and can't – do with three-wave panel data. *Sociol Method Res* 46: 44–67.
- Willett W, Rockström J, Loken B, Springmann M, Lan T, et al. (2019) Food in the Anthropocene: the EAT-Lancet Commission on healthy diets from sustainable food systems. *The Lancet*. DOI: 10.1016/S0140-6736(18)31788-4.
- World Bank (2018) *Climate Change Knowledge Portal*. [http://sdwebx.worldbank.org/climateportal/index.cfm?page=downscaled\\_data\\_download&menu=historical](http://sdwebx.worldbank.org/climateportal/index.cfm?page=downscaled_data_download&menu=historical). Accessed 28 May 2019.
- Young DJN, Stevens JT, Earles JM, Moore J, Ellis A, Jirka AL, Latimer AM (2017) Long-term climate and competition explain forest mortality patterns under extreme drought. *Ecol Lett* 20: 78-86.



## Appendices

### A. Methods and Materials

#### Global and regional forest carbon sink

To assess the net ecosystem exchange of carbon (NEE) of countries' forests I use the newest available direct measurements of NEE of 78 micrometeorological measurement towers located in forests of 16 countries from 2000-2014 provided by FLUXNET (NASA 2015). See Table B.1 in Appendix B for an overview of the analysed tower sites. FLUXNET sites collect data on the exchanges of CO<sub>2</sub> between forests and the atmosphere, precipitation and air temperature at least in a 30 minutes interval. Table B.3 provides a summary of the descriptive statistics. The tower sites use eddy covariance methods to measure forests' NEE. The unique dataset utilized here, 'FLUXNET2015', provides standardized values for these characteristics and underwent several quality control tests and gap-filling (Pastorello et al. 2017).

To infer the NEE of countries' forests from these 78 FLUXNET sites I apply a straightforward approach consisting of three steps: Firstly, I regress their annual NEE on several site characteristics (average temperature, average temperature squared, precipitation, latitude, and elevation) controlling for overall time-trends by including dummy variables of the years observed. While primarily interested in the variation between the forest sites, the inclusion of the 607 site-years available for this model minimize the influence of a specific observation period stemming from annual variation in climatic and other conditions. Therefore, all standard errors are clustered by tower site to ensure robustness with respect to heteroscedasticity and autocorrelation. The results of this linear ordinary least squares (OLS) regression model (Table B.2) indicate that only average temperature substantially relates to NEE. As Figure B.1 shows, the temperature – NEE relationship of forests follows a u-shaped pattern. Forests with an annual mean temperature of -5 to 0 °C are net emitters of carbon, whereas the carbon sequestration of forests is highest in climatic domains with an average of about 15 °C. Even higher temperatures are associated with lower sequestration. Note that uncertainty between 15 and 26 °C is relatively high, because of a rather limited number of tower sites in this climatic forest domain. The reported regression results of Table B.2 were tested for robustness: First, the model was rerun excluding one measurement tower each time from the regression. Second, all parameters were tested for linearity including a penalized splines fixed effects (FE) regression model (Ruppert et al. 2003). Furthermore, the robustness of standard errors was investigated via non-parametric bootstrapping. None of these

checks had any substantial influence on the estimates. In addition, the robustness of all estimates with respect to model specification was assessed using the procedure suggested by Young and Holsteen (2017). The potential influence of omitted variables was examined using the method suggested by Frank (2000). Also these checks detected no fundamental deviations from the reported results. The analyses were conducted using the statistical software package STATA 15.1.

Secondly, I predict the mean annual sequestration between the years 2000 and 2014 ( $t$ ) of country  $i$ 's forests in tons CO<sub>2</sub> per hectare ( $y_i$ ) from model 1 of Table B.2 according to the following formula:

$$y_i = \frac{1}{T} \sum_{t=1}^T (\beta_0 + \beta_1 a_{it} + \beta_2 a_{it}^2 + \beta_3 b_i + \gamma_t) \quad (\text{Eq. A.1}).$$

$\beta_0$  represents the model intercept.  $a_{it}$  stands for the average air temperature of country  $i$  in year  $t$ ,  $\beta_1$  for the regression coefficient of the sites' average temperature, and  $\beta_2$  for the coefficient of its square.  $b_i$  denotes country  $i$ 's centroid's latitude, and  $\beta_3$  the regression coefficient for the forest sites' latitude.  $\gamma_t$  represents the regression coefficient for year  $t$ . With  $\beta_0=5.60$ ,  $\beta_1=-2.20$ ,  $\beta_2=0.07$ ,  $\beta_3=-0.06$  from model 1 of Table B.2 follows:

$$y_i = \frac{1}{T} \sum_{t=1}^T (5.60 - 2.20a_{it} + 0.07a_{it}^2 - 0.06b_i + \beta_t) \quad (\text{Eq. A.2}).$$

Data for  $a_{it}$  is taken from the Climate Change Knowledge Portal of the World Bank (2018; Table B.6), and from the Country Geography Database of Portland State University (2018) for  $b_i$ . Computation of Eq. A.2 yields a global average of -8.8 tCO<sub>2</sub>ha<sup>-1</sup>yr<sup>-1</sup> (median = -9.2, sd. = 4.8, min. = -15.1, max. = 16.3) sequestered by forests in 2015. With roughly 2.7 trillion trees (Crowther et al. 2015) in the 40.0 Mkm<sup>2</sup> (FAO 2018) of forests worldwide, this translates into a mean of -8.8 kgCO<sub>2</sub>yr<sup>-1</sup> per tree (tropical forests (latitude 0° to <25° North (N) or South (S)): -8.4, temperate forests (25° to <50° N or S): -17.3, boreal forests (≥50° N): -1.9) as weighted by the share of trees by forest type (tropical: 0.48, temperate: 0.24, boreal: 0.27; Crowther et al. 2015).

Thirdly, simply multiplying countries' average NEE per hectare by their forest area gathered from the FAO (2018) gives countries' forest carbon sink. Summing up yields an estimate for the global forest carbon sequestration of -8.3 GtCO<sub>2</sub>yr<sup>-1</sup> or -1.1 tCO<sub>2</sub>yr<sup>-1</sup> per capita (p.c.; UNPD 2017).

## Afforestation/reforestation (AR) scenarios

To prevent dangerous climate change, the required and achievable forest land share of three different AR scenarios are developed. The basis for these scenarios is the 2 °C target and the associated remaining carbon budget until 2100. With the Paris Climate Agreement, the world community has agreed upon the limitation of global warming to well below 2 °C relative to preindustrial levels (UNFCCC 2015). A maximum of 2 °C of warming until 2100 may provide a relatively safe operating space for humanity and prevent dangerous climate change alongside a lock-in of a 'Hothouse Earth' pathway with potentially hazardous consequences for ecosystems and human socio-economic systems (IPCC 2014, Steffen et al. 2018, Fischer et al. 2018, Rockström et al. 2009). Yet, humanity allegedly has already committed to 1.3 °C of warming (Mauritsen and Pincus 2017). Hence, limiting global warming to 1.5 °C and presumably providing an even safer operating space (IPCC 2018) seems out of reach (Raftery et al. 2017). Global CO<sub>2</sub> emissions of fossil fuel use and industrial processes have risen to 35.8 GtCO<sub>2</sub> or 4.8 tCO<sub>2</sub> per capita (p.c.) in 2016 (Janssens-Maenhout et al. 2017). This surpasses the global annual gross carbon budget (an estimated 30 GtCO<sub>2</sub>) to fulfil the 2 °C target with a probability of at least 66 % (IPCC 2014, Friedlingstein et al. 2014, Meinshausen et al. 2009). Assuming an average annual world population of 9.8 billion people until 2100 (UNPD 2017) this goal translates into ~3 t of gross CO<sub>2</sub> emissions p.c. and year.

*Scenario 1* is the baseline assuming business-as-usual production and consumption patterns, constant other carbon sinks, further required emission reductions of 1.0 tCO<sub>2</sub>yr<sup>-1</sup> p.c. after accounting for the overall forest carbon sequestration of 0.8 tCO<sub>2</sub>yr<sup>-1</sup> p.c. with an expected average population of 9.8 billion people per year until 2100. Hence, the required additional absorption by forests for the 2 °C respectively the 3 t p.c. target is 9.8 GtCO<sub>2</sub>yr<sup>-1</sup>. Assuming similar carbon sequestration of established forests and afforested/reforested land, simple solution of the rule of three and addition to the existing forest area (40.0 Mkm<sup>2</sup>) delivers a required forest area of 88.0 Mkm<sup>2</sup>. With a global land area of 129.7 Mkm<sup>2</sup> (FAO 2018) this corresponds to a forest land share of 67.8 % necessary to reach the 2 °C target with AR activities alone. This implicitly assumes similar tree density, species, species richness and forest health of afforested/reforested land and established forests. To quantify the land area suitable for AR, land unsuitable for near-term and cost-efficient AR was excluded. These land cover types are artificial surfaces (including urban and associated areas), permanent snow and glaciers, terrestrial barren land, and sparsely natural vegetated areas as quantified by the FAO (2018). 100 % AR of all shrub-covered

areas and herbaceous vegetation (18.1 Mkm<sup>2</sup>) result in an achievable 44.8 % of forest land share in this scenario.

*Scenario 2* further assumes current diets and an associated demand of agricultural land of 2,100 m<sup>2</sup> p.c. (Hallström et al. 2015). As there are 3,770 m<sup>2</sup> p.c. of agricultural land currently available, 44 % of permanent grassland and cropland (FAO 2018) can be additionally afforested/reforested (16.4 Mkm<sup>2</sup>) for feeding an expected 9.8 billion people per year. Hence, a forest land share of 57.5 % can be realized in scenario 2 (77.6 Mkm<sup>2</sup>). This accounts for more than two thirds of the AR climate target outlined in scenario 1.

To achieve the AR target fully, further reduction in the demand for agricultural land is required. In *scenario 3* healthier diets with reduced red and ruminant meat consumption further decrease agricultural land demand by 28.0 % to 1,510 m<sup>2</sup> p.c. while dietary-related emissions decrease by 0.2 tCO<sub>2</sub>yr<sup>-1</sup> p.c. (Table 1 in Hallström et al. 2015). Hence, this reduction in carbon emissions implies a global reduction of the required carbon uptake by forests of 2.0 GtCO<sub>2</sub>yr<sup>-1</sup> to 7.8 GtCO<sub>2</sub>yr<sup>-1</sup>. This resembles a required forest land share of 60.0 % or 77.9 Mkm<sup>2</sup> of forest area. Via a further 28.0 % AR of permanent grassland and cropland a forest land share of 62.0 % or 80.4 Mkm<sup>2</sup> of forest area can be achieved to additionally sequester 8.0 GtCO<sub>2</sub>yr<sup>-1</sup>.

### Predictors of national forest land share

Compared to cross-sectional regression models, the FE panel model has the advantage of exploiting the longitudinal structure of the data as it only includes within-country variation. Hence, the FE model is not biased by cross-sectional unobserved heterogeneity (Brüderl and Ludwig 2015, Wooldridge 2010). If the strict exogeneity assumption ( $r(x_{it}, \varepsilon_{it}) = 0$ ) holds, FE models adequately estimate unbiased causal effects (Vaisey and Miles 2017). The model can be written as

$$y_{it} - \bar{y}_i = (x_{it} - \bar{x}_i)\beta + Z_t\gamma + \varepsilon_{it} - \bar{\varepsilon}_i \quad (\text{Eq. A.3}).$$

Here,  $y_{it}$  denotes the forest land share of country  $i$  in year  $t$ .  $\bar{y}_i$  represents country  $i$ 's mean of the whole observation period.  $x_{it}$  stands for the vector of all exogenous variables for country  $i$  at time  $t$ , and  $\bar{x}_i$  for the average of the time observed. The model further comprises a vector of dummy variables ( $Z$ ) for every year to control period effects for all countries (time FE). A country's time varying stochastic error term is represented by  $\varepsilon_{it}$ . All metric variables are included by taking their natural logarithm, which allows the estimation of elasticities. All standard errors are clustered by country and year, and are

therefore robust with respect to heteroscedasticity and autocorrelation. The reported regression results of Figure 4 were tested for robustness analogous to the results of the analysis for the FLUXNET data as already explained above. Furthermore, all six models were recalculated using the total forest land area as dependent variable instead of forest land share. None of these checks detected any substantial deviations from the results reported in Figure 4.

### **Additional references**

- Fischer H, Meissner KJ, Mix AC, Abram NJ, Austermann J, et al. (2018) Palaeoclimate constraints on the impact of 2 °C anthropogenic warming and beyond. *Nat Geosci* 11: 474-485.
- Frank KA (2000) Impact of a confounding variable on a regression coefficient. *Soc Method Res* 29: 147-194.
- IPCC – Intergovernmental Panel on Climate Change (2018) Global Warming of 1.5 °C. IPCC, Geneva.
- Mauritsen T, Pincus R (2017) Committed warming inferred from observations. *Nat Clim Change* 7: 652-655.
- Pastorello G, Papale D, Chu H, Trotta C, Agarwal DA, Canfora E, Baldocchi DD, Torn MS (2017) A new data set to keep a sharper eye on land-air exchanges. *Eos* 98. <https://doi.org/10.1029/2017EO071597>.
- Portland State University. Country Geography Database (2018). <https://www.pdx.edu/econ/country-geography-data>. Accessed 28 May 2019.
- Raftery AE, Zimmer A, Frierson DMW, Startz R, Liu P (2017) Less than 2 °C warming by 2100 unlikely. *Nat Clim Change* 7: 637-641.
- Rockström J, Steffen W, Noone K, Persson A, Chapin III FS, et al. (2009) A safe operating space for humanity. *Nature* 461: 472-475.
- Ruppert D, Wand MP, Carroll RJ (2003) Semiparametric regression. Cambridge University Press, Cambridge, UK.
- Steffen W, Rockström J, Richardson K, Lenton TM, Folke C, et al. (2018) Trajectories of the earth system in the anthropocene. *P Natl Acad Sci USA* 115: 8252-8259.
- UNFCCC – United Nations Framework Convention on Climate Change (2015) Adoption of the Paris Agreement Report No. FCCC/CP/2015/L.9/Rev.1.
- Wooldridge JM (2010) Econometric analysis of cross-section and panel data. MIT Press, Cambridge.
- Young C, Holsteen K (2017) Model uncertainty and robustness: A computational framework for multimodel analysis. *Soc Method Res* 46: 3-40.

**Data availability:** All data used in this article is publicly available at the referenced webpages.


## B. Supplementary Figures and Tables

Country	Tower site code	Reference paper	Country	Tower site code	Reference Paper	Country	Tower site code	Reference paper
Belgium	BE-Vie	Aubinet et al. (2001)	French	GF-Guy	Bonal et al. (2008)	United States	US-Blo	Falge et al. (2002)
Brazil	BR-Sa1	Hayek et al. (2018)	Guyana				US-GBT	Zeller and Nikolov (2000)
	BR-Sa3	Saleska et al. (2003)	Germany	DE-Hai	Knohl et al. (2003)		US-GLF	Frank et al. (2014)
Canada	CA-Gro	McCaughey et al. (2006)		DE-Lkb	Lindauer et al. (2014)		US-Ha1	Barford et al. (2001)
	CA-NS1	Goulden et al. (2006)		DE-Lnf	Anthoni et al. (2004)		US-KS1	Dore et al. (2003)
	CA-NS2	Bond-Lamberty et al. (2004)		DE-Obe	Bernhofer et al. (2008)		US-Me1	Irvine et al. (2007)
	CA-NS3	Bond-Lamberty et al. (2004)	Ghana	DE-Tha	Grünwald and Bernhofer (2007)		US-Me2	Law et al. (2004)
	CA-NS4	Schmidt et al. (2011)		GH-Ank	Chiti et al. (2010)		US-Me3	Sun et al. (2004)
	CA-NS5	Bond-Lamberty et al. (2004)	Italy	IT-CA1	Sabbatini et al. (2016)		US-Me4	Law et al. (2004)
	CA-Oas	Chen et al. (2003)		IT-CA3	Sabbatini et al. (2016)		US-Me5	Law et al. (2004)
	CA-Obs	Chen et al. (2003)		IT-Cp2	Fares et al. (2014)		US-Me6	Ruehr et al. (2012)
	CA-Qfo	Bergeron et al. (2007)		IT-Cpz	Garbulsky et al. (2008)		US-MMS	Baldocchi et al. (2005)
	CA-SF1	Amiro et al. (2006)		IT-Isp	Ferréa et al. (2012)		US-Oho	Chu et al. (2016)
	CA-TP1	Arain and Restrepo-Coupe (2005)		IT-La2	Marcolla et al. (2003)		US-PFa	Desai et al. (2008)
	CA-TP2	Arain and Restrepo-Coupe (2005)		IT-Lav	Marcolla et al. (2003)		US-Prr	Kobayashi et al. (2014)
	CA-TP3	Arain and Restrepo-Coupe (2005)		IT-PT1	Migliavacca et al. (2009)		US-Syv	Desai et al. (2005)
	CA-TP4	Arain and Restrepo-Coupe (2005)		IT-Ren	Montagnani et al. (2009)		US-UMd	Gough et al. (2013)
	CA-TPD	Schmidt et al. (2011)		IT-Ro1	Rey et al. (2002)		US-WCr	Desai et al. (2005)
Czech Republic	CZ-BK1	Acosta et al. (2013)		IT-Ro2	Tedeschi et al. (2006)		US-Wi0	Desai et al. (2008)
Denmark	DK-Sor	Pilegaard et al. (2011)		IT-SR2	Gruening et al. (2013)		US-Wi1	Desai et al. (2008)
Finland	FI-Hyy	Suni et al. (2003)	Netherlands	IT-SRo	Chiesi et al. (2005)		US-Wi2	Desai et al. (2008)
	FI-Let	Koskinen et al. (2014)		NL-Loo	Moors (2012)		US-Wi3	Desai et al. (2008)
	FI-Sod	Thum et al. (2007)	Panama	PA-SPn	Wolf et al. (2011)		US-Wi4	Desai et al. (2008)
France	FR-Fon	Delpierre et al. (2016)	Russian Federation	RU-Fyo	Kurbatova et al. (2008)		US-Wi5	Schmidt et al. (2011)
	FR-LBr	Berbigier et al. (2001)		RU-SkP	Maximov (2012)		US-Wi8	Desai et al. (2008)
	FR-Pue	Rambal et al. (2004)	Switzerland	CH-Dav	Zielis et al. (2014)		US-Wi9	Schmidt et al. (2011)
				CH-Lae	Etzold et al. (2011)			

**Table B.1 | FLUXNET micrometeorological measurement towers in forests included in the regression analysis by country (NASA 2015).**

Model	(1)
Dependent variable	NEE
Average temperature	-2.20*** (0.46)
Average temperature squared	0.07** (0.03)
Precipitation	-0.00 (0.00)
Latitude	-0.06 (0.29)
Elevation	-0.00 (0.00)
2000 (reference)	
2001	3.80 (2.60)
2002	-1.21 (3.38)
2003	-1.77 (3.98)
2004	-2.15 (3.69)
2005	-1.42 (3.65)
2006	-0.10 (3.62)
2007	-3.53 (3.84)
2008	-1.75 (3.73)
2009	-1.45 (4.02)
2010	0.34 (3.83)
2011	-1.63 (3.90)
2012	-1.24 (3.90)
2013	-0.93 (3.80)
2014	-2.46 (3.92)
Constant	5.60 (19.49)
n x T	607
n	78
adjusted R <sup>2</sup>	0.15

**Table B.2 | Linear OLS regression of net ecosystem exchange.** NEE = net ecosystem exchange in  $\text{tCO}_2\text{ha}^{-1}\text{yr}^{-1}$ . \*\* =  $p < 0.01$ , \*\*\* =  $p < 0.001$ . Unstandardized regression coefficients with standard errors in brackets. All standard errors are clustered by tower site, and robust with respect to heteroscedasticity and autocorrelation. Years covered: 2000-2014. Table B.3 gives a descriptive overview of all variables in model 1. Table B.1 lists all 78 micrometeorological measurement towers of FLUXNET in forests included in model 1.



**Figure B.1 | Predicted values of net ecosystem exchange (NEE) by annual average temperature.** NEE of CO<sub>2</sub> as predicted by the OLS regression model presented in Table B.2 (dark blue line) with 95% confidence intervals (blue area). Negative numbers on the y-axis indicate net absorption of CO<sub>2</sub> by forests, positive numbers net CO<sub>2</sub> release.

Variable	mean	within ( $\bar{x}_i$ )		between ( $x_{it} - \bar{x}_i + \bar{x}$ )			N (nXT)	n	Description	
		sd	min.	max.	sd	min.				max.
Net ecosystem exchange	-13.98	6.28	-41.86	13.55	15.52	-67.40	10.21	674	94	Net ecosystem exchange (NEE) of CO <sub>2</sub> . NEE is the sum of Gross Primary Productivity (GPP, i.e. biomass stored) and ecosystem respiration (release of CO <sub>2</sub> from soil and plant). Negative numbers indicate net absorption, positive numbers net release of CO <sub>2</sub> . Unit: t per ha.
Average temperature	8.87	0.73	6.52	11.21	7.42	-4.62	25.89	693	94	Average annual air temperature derived from daily averages. Unit: °C.
Precipitation	0.92	0.19	0.19	1.59	0.56	0.16	3.11	693	94	Annual precipitation. Sum of daily data. Unit: 1000 mm.
Latitude	44.39	0	44.39	44.39	12.93	2.9	67.4	693	94	In degrees north or south from equator.
Elevation	527.62	0	527.62	527.62	596.38	1	3197	625	78	Elevation of site. Unit: m above sea level.

**Table B.3 | Variable description of FLUXNET data of micrometeorological measurement towers in forests.** Data source is FLUXNET, a global network of micrometeorological tower sites with long-term measurement. FLUXNET is operated by the Oak Ridge National Laboratory Distributed Active Archive Center (ORNL DAAC) of the National Aeronautics and Space Administration (NASA) of the United States. Years covered: 2000-2014.



Rank	Country	Sum score of quintiles	CO <sub>2</sub> emissions	CO <sub>2</sub> p.c. emissions	Forest land share	NEE per ha	NEE	NEE potential	GDP p.c.
1	Japan	33	5	5	5	4	5	4	5
2	Spain	33	5	4	4	5	5	5	5
3	South Korea	32	5	5	5	5	4	3	5
4	France	32	5	4	3	5	5	5	5
5	United States	31	5	5	3	3	5	5	5
6	Australia	31	5	5	2	4	5	5	5
7	Mexico	30	5	4	3	4	5	5	4
8	Argentina	30	5	4	2	5	5	5	4
9	Italy	30	5	4	3	5	4	4	5
10	Germany	30	5	5	3	4	4	4	5
11	Turkey	30	5	4	2	5	5	5	4
12	Brazil	29	5	3	5	2	5	5	4
13	Peru	29	4	3	5	4	5	5	3
14	New Zealand	29	3	5	4	4	4	4	5
15	China	29	5	5	3	3	5	5	3
16	Poland	29	5	5	3	4	4	4	4
17	Iran, Islamic Rep.	29	5	5	1	5	4	5	4
18	Venezuela, RB	29	5	4	5	2	5	4	4
19	South Africa	28	5	5	1	5	4	5	3
20	Turkmenistan	28	4	5	2	5	4	4	4
21	Malaysia	28	5	5	5	2	4	3	4
22	Belarus	28	4	4	4	4	4	4	4
23	Czech Republic	28	4	5	4	4	3	3	5
24	Greece	28	4	4	3	5	4	4	4
25	Bulgaria	28	4	5	4	5	3	3	4
26	United Kingdom	27	5	4	2	4	3	4	5
27	Portugal	27	4	4	4	5	3	3	4
28	Chile	27	4	4	3	3	5	4	4
29	Serbia	27	5	5	3	5	3	3	3
30	Bolivia	27	3	3	5	4	5	5	2
31	Romania	27	4	4	3	4	4	4	4
32	Indonesia	26	5	3	5	1	5	4	3
33	Finland	26	4	5	5	1	4	2	5
34	Austria	26	4	5	4	3	3	2	5
35	Hungary	26	4	4	3	5	3	3	4
36	Belgium	26	4	5	3	5	2	2	5
37	Kazakhstan	26	5	5	1	3	3	5	4
38	Ukraine	26	5	4	2	4	4	5	2
39	Colombia	26	4	3	5	2	5	4	3
40	Slovenia	25	3	5	5	4	2	1	5
41	Thailand	25	5	4	3	1	4	4	4
42	Croatia	25	3	4	4	5	3	2	4
43	India	25	5	3	3	2	5	5	2
44	Vietnam	25	5	3	5	2	4	4	2
45	Sweden	25	4	4	5	1	4	2	5
46	Morocco	25	4	3	2	5	4	4	3
47	Slovak Republic	25	3	5	4	4	2	2	5
48	Bosnia and Herzegovina	24	3	5	4	4	3	2	3
49	Angola	24	3	2	4	3	5	5	2
50	Netherlands	24	4	5	2	5	1	2	5
51	Ecuador	24	3	3	5	3	4	3	3
52	Ireland	24	3	5	2	4	2	3	5
53	Estonia	24	3	5	5	3	2	1	5
54	Latvia	23	2	4	5	3	3	2	4
55	Zambia	23	2	1	5	3	5	5	2
56	Zimbabwe	23	3	2	4	3	5	5	1
57	Norway	23	4	5	3	1	3	2	5
58	Botswana	23	2	3	2	3	4	5	4
59	Uruguay	23	2	3	2	5	3	4	4
60	Azerbaijan	23	3	4	2	5	2	3	4
61	Myanmar	23	3	1	4	3	5	5	2
62	Uzbekistan	22	4	3	1	5	3	4	2
63	Paraguay	22	2	2	4	3	4	4	3
64	Russian Federation	22	5	5	5	1	1	1	4
65	Namibia	22	1	3	2	4	4	5	3
66	Lithuania	22	3	4	4	3	2	2	4
67	Syrian Arab Republic	22	4	3	1	5	1	3	5
68	Denmark	22	3	4	2	4	2	2	5
69	Lao PDR	22	1	2	5	3	5	4	2
70	Canada	22	5	5	4	1	1	1	5
71	Tanzania	22	2	1	5	3	5	5	1
72	Iraq	21	4	4	1	4	2	3	3


**Table B.4 | Full country ranking of climate responsibility, forests' mitigation contribution and potential, and economic capabilities in 2015.**

Rank	Country	Sum score of quintiles	CO <sub>2</sub> emissions	CO <sub>2</sub> p.c. emissions	Forest land share	NEE per ha	NEE	NEE potential	GDP p.c.
73	Congo, Rep.	21	2	2	5	2	5	3	2
74	Israel	21	4	5	1	4	1	1	5
75	Switzerland	21	3	4	3	3	2	1	5
76	Gabon	21	2	3	5	2	4	1	4
77	Algeria	21	4	4	1	3	2	4	3
78	Congo, Dem. Rep.	20	2	1	5	2	5	4	1
79	Cameroon	20	3	2	4	2	4	3	2
80	Panama	20	2	3	5	2	3	1	4
81	Georgia	20	2	3	4	3	3	2	3
82	Lebanon	20	3	4	2	5	1	1	4
83	Saudi Arabia	20	5	5	1	2	1	1	5
84	Pakistan	20	4	2	1	4	2	5	2
85	Central African Republic	20	1	1	4	2	4	3	5
86	Macedonia, FYR	20	2	4	4	4	2	1	3
87	Cuba	20	3	3	3	2	2	2	5
88	Papua New Guinea	20	2	2	5	2	5	2	2
89	Mozambique	20	2	1	5	2	5	4	1
90	Albania	19	1	3	3	5	2	2	3
91	Dominican Republic	19	3	3	4	3	2	1	3
92	Tunisia	19	3	3	1	4	2	3	3
93	Philippines	19	4	2	3	2	3	3	2
94	Egypt, Arab Rep.	19	5	3	1	3	1	3	3
95	Nepal	18	2	1	3	5	3	3	1
96	Ethiopia	18	2	1	2	3	4	5	1
97	Costa Rica	18	2	3	5	2	2	1	3
98	Libya	18	4	5	1	3	1	2	2
99	Honduras	17	2	2	4	2	3	2	2
100	Bangladesh	17	4	2	2	2	2	3	2
101	Nigeria	17	4	2	1	1	3	4	2
102	Kenya	17	3	2	1	2	3	5	1
103	Ghana	17	3	2	4	1	3	2	2
104	Afghanistan	17	2	1	1	5	2	5	1
105	Cambodia	17	2	2	5	1	3	2	2
106	Somalia	17	1	1	2	1	3	4	5
107	Guatemala	17	3	2	3	2	3	2	2
108	Moldova	17	2	3	2	5	1	2	2
109	Cote d'Ivoire	16	2	2	3	1	3	3	2
110	Jordan	16	3	3	1	4	1	1	3
111	Swaziland	16	1	2	4	4	1	1	3
112	Madagascar	16	1	1	2	3	4	4	1
113	Jamaica	15	2	3	3	2	1	1	3
114	Malawi	15	1	1	3	3	3	3	1
115	Sri Lanka	15	3	2	3	1	2	1	3
116	Nicaragua	15	2	2	3	2	2	2	2
117	Uganda	14	2	1	2	2	2	4	1
118	Mongolia	14	3	4	1	1	1	1	3
119	Burundi	14	1	1	2	3	1	1	5
120	Liberia	14	1	1	4	2	3	2	1
121	Senegal	14	2	2	4	1	2	2	1
122	Armenia	14	1	3	2	3	1	1	3
123	Yemen, Rep.	13	3	2	1	3	1	2	1
124	Lesotho	13	1	1	1	5	1	2	2
125	Guinea	13	1	1	3	1	3	3	1
126	Benin	13	2	2	4	1	2	1	1
127	El Salvador	13	2	2	2	2	1	1	3
128	Kyrgyz Republic	12	2	2	1	1	1	3	2
129	Rwanda	11	1	1	2	4	1	1	1
130	Guinea-Bissau	11	1	1	5	1	1	1	1
131	Chad	11	1	1	1	1	2	4	1
132	Sierra Leone	11	1	1	4	1	2	1	1
133	Tajikistan	11	1	2	1	2	1	3	1
134	Gambia, The	11	1	1	5	1	1	1	1
135	Burkina Faso	10	1	1	2	1	2	2	1
136	Mauritania	10	1	2	1	1	1	2	2
137	Niger	9	1	1	1	1	1	3	1
138	Mali	9	1	1	1	1	1	3	1
139	Haiti	8	1	1	1	2	1	1	1
140	Eritrea	8	1	1	2	1	1	1	1
141	Togo	7	1	1	1	1	1	1	1

**Table B.4, continued | Full country ranking of climate responsibility, forests' mitigation contribution and potential, and economic capabilities in 2015.** p.c. = per capita, NEE = net ecosystem exchange, GDP = gross domestic product. Numbers represent the quintiles the countries rank if not indicated otherwise. Data sources: CO<sub>2</sub> emissions: EDGAR – Emissions Database for Global Atmospheric Research; forest land share: FAO – Food and Agriculture Organization of the UN; NEE: own calculations based on FLUXNET data; GDP: IMF – International Monetary Fund.

Model	(1)	(2)	(3)	(4)	(5)	(6)
Dependent variable	Forest land share					
Agricultural land	-0.21* (0.10)	-0.08 (0.10)	-0.03 (0.10)	-0.04 (0.11)	0.05 (0.13)	0.03 (0.12)
Population		-0.27** (0.08)	-0.28** (0.08)	-0.27** (0.08)	-0.17* (0.08)	-0.18* (0.08)
Urban population		-0.00 (0.09)	-0.02 (0.09)	-0.01 (0.09)	-0.12 (0.09)	-0.11 (0.08)
GDP per capita			0.10* (0.04)	0.09* (0.03)	0.10* (0.04)	0.09* (0.04)
Industry			-0.10* (0.04)	-0.08* (0.04)	-0.09* (0.04)	-0.10* (0.04)
Services			-0.08 (0.05)	-0.08+ (0.04)	-0.09+ (0.05)	-0.09+ (0.05)
Forest products trade balance				0.02 (0.03)	0.02 (0.02)	0.03 (0.02)
Protected forest area					0.06* (0.03)	0.06* (0.03)
Mean temperature						-0.10* (0.04)
Droughts						0.00 (0.00)
n x T	2494	2494	2255	2255	1781	1744
n	98	98	96	96	88	88
adjusted R <sup>2</sup> within	0.06	0.15	0.22	0.20	0.27	0.28

**Table B.5 | Country and time fixed effects regressions of forest land share.** + =  $p < 0.10$ , \* =  $p < 0.05$ , \*\* =  $p < 0.01$ . Unstandardized regression coefficients with standard errors in brackets. All six models include the years 1990-2015 and contain dummy variables for each year in order to control for overall time-trends. All standard errors are clustered by country and year, and robust with respect to heteroscedasticity and autocorrelation.



**Figure B.2 | Partial residual plot for GDP p.c. of model 6 in Figure 4 and Table B.5.** Partial residual for every country year (blue filled circles) and the smoothed mean (red curve) as calculated from the fixed effects regression with penalized splines (Ruppert et al. 2003) for logged GDP per capita. Red ticks on the x-axis represent knots. The plot demonstrates that the effect of GDP growth on forest land share growth is almost flat for poor countries with logged PPP GDP p.c. of less than ca. 8.0, and positive and virtually linear for richer countries. Thus, the effect is positive and linear for the vast majority of observations.

Variable	mean	within ( $\bar{x}_i$ )			between ( $x_{it} - \bar{x}_i + \bar{x}$ )			N	n	Description	Data Source
		sd	min.	max.	sd	min.	max.				
Forest land share	32.77	2.02	18.34	50.01	23.83	0	98.60	4690	181	Forest is determined both by the FAO presence of trees and the absence of other predominant land uses. Forest area is land under natural or planted stands of trees of at least 5 meters in situ or with the potential of growth to this height, with an area of more than 0.5ha and width of more than 20m, and a canopy cover of at least 10%, whether productive or not, and excludes tree stands in agricultural production systems and trees in urban parks and gardens. Unit: % of land area.	
Agricultural land	40.37	2.87	19.71	57.93	21.12	0.53	84.40	4650	182	Agricultural land refers to the share FAO of land area that is arable, under permanent crops, and under permanent pastures. Unit: % of land area.	
Population	34.12	12.17	-191.34	249.11	127.76	0.02	1272.37	4754	183	Total population. Unit: 1 million.	UNPD, WB
Urban population	52.83	3.37	39.20	69.73	23.32	8.93	100	4806	185	Urban population refers to people living in urban areas as defined by FAO, national statistical offices. Unit: % of total population.	WB
GDP p. c.	11.29	4.85	-16.87	47.98	11.75	0.54	62.31	4384	177	Gross domestic product (GDP) per IMF capita (p.c.) based on purchasing power parity (PPP). PPP GDP is GDP converted to international dollars using PPP rates. Unit: 1000 international dollars.	
Industry, value added	28.17	4.50	-5.59	60.52	11.04	7.20	75.96	4092	174	Industry corresponds to the WB International Standard Industrial Classification (ISIC) divisions 10-45. The origin of value added is determined by the ISIC, revision 3. Unit: % of GDP.	
Services, value added	55.57	5.41	22.75	99.17	13.46	22.77	82.07	4075	173	Services correspond to ISIC WB divisions 50-99. The industrial origin of value added is determined by the ISIC, revision 3. Unit: % of GDP.	
Forest products trade balance	0.16	1.53	-24.07	27.69	2.03	-2.09	24.27	4171	174	Forest products trade balance is the FAO ratio of exports to imports of forest goods as share of GDP.	
Protected forest area	37.40	34.56	-640.36	469.28	142.59	0	1630.39	3289	149	Protected forest area is designated FAO primarily for conservation of biological diversity and natural and associated cultural resources. Protection and maintenance is managed through legal or other effective means. Unit: km <sup>2</sup> .	
Mean temperature	18.98	0.49	15.61	21.29	8.21	-5.97	28.90	4225	169	Mean annual air temperature WB derived from quality controlled monthly observational data from thousands of weather stations worldwide. Unit: °C.	
Droughts	0.20	0.36	-0.60	1.16	0.24	0	1	1963	160	Dummy, 1, if a drought occurred at least once a year. A drought is classified if at least one of the following criteria is met: 10 or more people dead, 100 or more people affected, declaration of a state of emergency, call for international assistance.	

**Table B.6 | Drivers of national forest land share: variable description.** CRED = Centre for Research on the Epidemiology of Disasters, FAO = Food and Agriculture Organization of the United Nations, IMF = International Monetary Fund, UNPD = United Nations Population Division, WB = World Bank; All variables in the models are included by taking the natural logarithm allowing for the estimation of elasticities. Years covered: 1990-2015.

---

Algeria**	Hungary**	Peru**
Argentina**	India**	Philippines**
Australia**	Indonesia**	Poland**
Austria**	Iran, Islamic Rep.**	Portugal**
Bangladesh**	Ireland**	Romania**
Belarus**	Israel	Russian Federation**
Belgium**	Italy**	Senegal**
Brazil**	Jamaica**	Serbia**
Bulgaria**	Japan*	Slovak Republic**
Burkina Faso**	Kenya**	Slovenia**
Cambodia**	Lao PDR*	South Africa**
Cameroon**	Latvia**	South Korea**
Canada**	Lebanon**	Spain**
Chile**	Liberia	Sri Lanka*
China**	Lithuania**	Swaziland**
Colombia**	Malawi**	Sweden**
Congo, Rep.**	Malaysia**	Switzerland**
Costa Rica*	Mali**	Tajikistan**
Croatia**	Mexico**	Tanzania**
Czech Republic**	Mongolia**	Thailand**
Denmark**	Morocco**	Tunisia**
Dominican Republic*	Mozambique**	Turkey**
Ecuador**	Myanmar**	Uganda**
Estonia**	Namibia**	Ukraine**
Ethiopia*	Nepal**	United Kingdom*
Finland**	Netherlands**	United States**
France*	New Zealand**	Uruguay**
Gabon**	Nicaragua**	Uzbekistan**
Gambia, The**	Niger**	Venezuela, RB**
Georgia**	Norway**	Vietnam**
Germany**	Panama**	Zambia**
Ghana**	Papua New Guinea**	Zimbabwe**
Guatemala**	Paraguay**	

---

**Table B.7 | Countries included in the analyses.** All 98 countries are full members of the United Nations, have sufficient quality of forest area data (tier 2 and 3; FAO 2016) and are included in the models 1, and 2 of Table B.5. Due to missing values in the further added variables, the models 3 and 4 include the 96 countries indicated by '\*', and for the models 5 and 6 of Table B.5 the 88 countries marked by '#'. .

## Additional references

- Acosta M, Pavelka M, Montagnani L, Kutsch W, Lindroth A, Juszczak R, Janouš D (2013) Soil surface CO<sub>2</sub> efflux measurements in Norway spruce forests: Comparison between four different sites across Europe — from boreal to alpine forest. *Geoderma* 192: 295–303.
- Amiro B, Barr A, Black T, Iwashita H, Kljun N, McCaughey J, Morgenstern K, Murayama S, Nesic Z, Orchansky A (2006) Carbon, Energy And Water Fluxes At Mature And Disturbed Forest Sites, Saskatchewan, Canada. *Agric For Meteorol* 136: 237-251.
- Anthoni PM, Knohl A, Rebmann C, Freibauer A, Mund M, Ziegler W, Kolle O, Schulze E-D (2004) Forest and agricultural land-use-dependent CO<sub>2</sub> exchange in Thuringia, Germany. *Glob Chang Biol* 10: 2005–2019.
- Arain MA, Restrepo-Coupe N (2005) Net Ecosystem Production In A Temperate Pine Plantation In Southeastern Canada. *Agric For Meteorol* 128: 223-241.
- Aubinet M, Chermanne B, Vandenhaute M, Longdoz B, Yernaux M, Laitat E (2001) Long term carbon dioxide exchange above a mixed forest in the Belgian Ardennes. *Agric For Meteorol* 108: 293–315.
- Baldocchi DD, Black TA, Curtis PS, Falge E, Fuentes JD, Granier A, Gu L, Knohl A, Lee X, Pilegaard K, Schmid HP, Valentini R, Wilson K, Wofsy S, Xu L, Yamamoto S (2005) Predicting The Onset Of Net Carbon Uptake By Deciduous Forests With Soil Temperature And Climate Data: A Synthesis Of FLUXNET Data. *Int J Biometeorol*: 49: 377-387.
- Barford CC, Wofsy SC, Goulden ML, Munger JW, Pyle EH, Urbanski SP, Hutyra L, Saleska SR, Fitzjarrald D, Moore K (2001) Factors Controlling Long- And Short-Term Sequestration Of Atmospheric CO<sub>2</sub> In A Mid-Latitude Forest. *Science* 294: 1688-1691.
- Berbigier P, Bonnefond J-M, Mellmann P (2001) CO<sub>2</sub> and water vapour fluxes for 2 years above Euroflux forest site. *Agric For Meteorol* 108: 183–197.
- Bergeron O, Margolis HA, Black TA, Coursolle C, Dunn AL, Barr AG, Wofsy SC (2007) Comparison Of Carbon Dioxide Fluxes Over Three Boreal Black Spruce Forests In Canada. *Glob Chang Biol* 13: 89-107.
- Bernhofer C, Grünwald T, Moderow U, Hehn M, Eichelmann U, Prasse H (2008) FLUXNET2015 DE-Obe Oberbärenburg. DOI: 10.18140/FLX/1440151.
- Bonal D, Bosc A, Ponton S, Goret J-Y, Burban B, Gross P, Bonnefond J-M, Elbers J, Longdoz B, Epron D, Guehl J-M, Granier A (2008) Impact of severe dry season on net ecosystem exchange in the Neotropical rainforest of French Guiana. *Glob Chang Biol* 14: 1917–1933.
- Bond-Lamberty B, Wang C, Gower ST (2004) Net Primary Production And Net Ecosystem Production Of A Boreal Black Spruce Wildfire Chronosequence. *Glob Chang Biol* 10: 473-487.
- Chen JM, Ju W, Cihlar J, Price D, Liu J, Chen W, Pan J, Black A, Barr A (2003) Spatial Distribution Of Carbon Sources And Sinks In Canadas Forests. *Tellus B* 55: 622-641.
- Chiesi M, Maselli F, Bindi M, Fibbi L, Cherubini P, Arlotta E, Tirone G, Matteucci G, Seufert G (2005) Modelling carbon budget of Mediterranean forests using ground and remote sensing measurements. *Agric For Meteorol* 135: 22–34.
- Chiti T, Certini G, Grieco E, Valentini R (2010) The role of soil in storing carbon in tropical rainforests: the case of Ankasa Park, Ghana. *Plant Soil* 331: 453-461.
- Chu H, Chen J, Gottgens JF, Desai AR, Ouyang Z, Qian SS (2016) Response And Biophysical Regulation Of Carbon Dioxide Fluxes To Climate Variability And Anomaly In Contrasting Ecosystems In Northwestern Ohio, USA. *Agric For Meteorol* 220: 50-68.
- Delpierre N, Berveiller D, Granda E, Dufrêne E (2016) Wood phenology, not carbon input, controls the interannual variability of wood growth in a temperate oak forest. *New Phytol* 210: 459-470.

- Desai AR, Bolstad PV, Cook BD, Davis KJ, Carey EV (2005) Comparing Net Ecosystem Exchange Of Carbon Dioxide Between An Old-Growth And Mature Forest In The Upper Midwest, USA. *Agric For Meteorol* 128: 33-55.
- Desai AR, Noormets A, Bolstad PV, Chen J, Cook BD, Davis KJ, Euskirchen ES, Gough C, Martin JG, Ricciuto DM, Schmid HP, Tang J, Wang W (2008) Influence Of Vegetation And Seasonal Forcing On Carbon Dioxide Fluxes Across The Upper Midwest, USA: Implications For Regional Scaling. *Agric For Meteorol* 148: 288-308.
- Dore S, Hymus GJ, Johnson DP, Hinkle CR, Valentini R, Drake BG (2003) Cross Validation Of Open-Top Chamber And Eddy Covariance Measurements Of Ecosystem CO<sub>2</sub> Exchange In A Florida Scrub-Oak Ecosystem. *Glob Chang Biol* 9: 84-95.
- Etzold S, Ruehr NK, Zweifel R, Dobbertin M, Zingg A, Pluess P, Häsler R, Eugster W, Buchmann N (2011) The Carbon Balance of Two Contrasting Mountain Forest Ecosystems in Switzerland: Similar Annual Trends, but Seasonal Differences. *Ecosystems* 14: 1289–1309.
- Falge E, Baldocchi D, Tenhunen J, Aubinet M, Bakwin P, Berbigier P, Bernhofer C, Burba G, Clement R, Davis KJ, Elbers JA, Goldstein AH, Grelle A, Granier A, Guðmundsson J, Hollinger D, Kowalski AS, Katul G, Law BE, Malhi Y, Meyers T, Monson RK, Munger J, Oechel W, Paw UKT, Pilegaard K, Rannik Ü, Rebmann C, Suyker A, Valentini R, Wilson K, Wofsy S (2002) Seasonality Of Ecosystem Respiration And Gross Primary Production As Derived From FLUXNET Measurements. *Agric For Meteorol* 113: 53-74.
- Fares S, Savi F, Muller J, Matteucci G, Paoletti E (2014) Simultaneous measurements of above and below canopy ozone fluxes help partitioning ozone deposition between its various sinks in a Mediterranean Oak Forest. *Agric For Meteorol* 198: 181–191.
- Ferréa C, Zenone T, Comolli R, Seufert G (2012) Estimating heterotrophic and autotrophic soil respiration in a semi-natural forest of Lombardy, Italy. *Pedobiologia* 55: 285-294.
- Frank JM, Massman WJ, Ewers BE, Huckaby LS, Negrón JF (2014) Ecosystem CO<sub>2</sub> /H<sub>2</sub>O Fluxes Are Explained By Hydraulically Limited Gas Exchange During Tree Mortality From Spruce Bark Beetles. *Journal of Geophysical Research: Biogeosciences* 119: 1195-1215.
- Garbulsky MF, Penuelas J, Papale D, Filella I (2008) Remote estimation of carbon dioxide uptake by a Mediterranean forest. *Glob Chang Biol* 14: 2860–2867.
- Gough CM, Hardiman BS, Nave LE, Bohrer G, Maurer KD, Vogel CS, Nadelhoffer KJ, Curtis PS (2013) Sustained Carbon Uptake And Storage Following Moderate Disturbance In A Great Lakes Forest. *Ecol Appl* 23: 1202-1215.
- Goulden ML, Winston GC, McMillan AM, Litvak ME, Read EL, Rocha AV, Rob Elliot J (2006) An Eddy Covariance Mesonet To Measure The Effect Of Forest Age On Land-Atmosphere Exchange. *Glob Chang Biol* 12: 2146-2162.
- Gruening C, Godeed I, Cescatti A, Pokorska O (2013) FLUXNET2015 IT-SR2 San Rossore 2. DOI: 10.18140/FLX/1440236.
- Grünwald T, Bernhofer C (2007) A decade of carbon, water and energy flux measurements of an old spruce forest at the Anchor Station Tharandt. *Tellus B* 59: 387–396.
- Hayek MN, Wehr R, Longo M, Hutrya LR, Wiedemann K, Munger JW, Bonal D, Saleska SR, Fitzjarrald DR, Wofsy SC (2018) A Novel Correction For Biases In Forest Eddy Covariance Carbon Balance. *Agric Forest Meteorol* 250: 90-101.
- Irvine J, Law BE, Hibbard KA (2007) Postfire Carbon Pools And Fluxes In Semiarid Ponderosa Pine In Central Oregon. *Glob Chang Biol* 13: 1748-1760.
- Knohl A, Schulze E-D, Kolle O, Buchmann N (2003) Large carbon uptake by an unmanaged 250-year-old deciduous forest in Central Germany. *Agric For Meteorol* 118: 151–167.



- Kobayashi H, Suzuki R, Nagai S, Nakai T, Kim Y (2014) Spatial Scale And Landscape Heterogeneity Effects On Fapar In An Open-Canopy Black Spruce Forest In Interior Alaska. *IEEE Geosci Remote S* 11: 564-568.
- Koskinen M, Minkkinen K, Ojanen P, Kämäräinen M, Laurila T, Lohila A (2014) Measurements of the CO<sub>2</sub> exchange with an automated chamber system throughout the year: challenges in measuring the nighttime respiration in porous peat soil. *Biogeosciences* 11: 347-363.
- Kurbatova J, Li C, Varlagin A, Xiao X, Vygodskaya N (2008) Modeling carbon dynamics in two adjacent spruce forests with different soil conditions in Russia. *Biogeosciences* 5: 969–980.
- Law BE, Turner D, Campbell J, Sun OJ, Van Tuyl S, Ritts WD, Cohen WB (2004) Disturbance And Climate Effects On Carbon Stocks And Fluxes Across Western Oregon USA. *Glob Chang Biol* 10: 1429-1444.
- Lindauer M, Schmid HP, Grote R, Mauder M, Steinbrecher R, Wolpert B (2014) Net ecosystem exchange over a non-cleared wind-throw-disturbed upland spruce forest—Measurements and simulations. *Agric For Meteorol* 197: 219–234.
- Marcolla B, Pitacco A, Cescatti A (2003) Canopy architecture and turbulence structure in a Coniferous forest. *Bound-Lay Meteorol* 108: 39–59.
- Maximov T (2012) FLUXNET2015 RU-SkP Yakutsk Spasskaya Pad larch. DOI:10.18140/FLX/1440243.
- McCaughey J, Pejam M, Arain M, Cameron D (2006) Carbon Dioxide And Energy Fluxes From A Boreal Mixedwood Forest Ecosystem In Ontario, Canada. *Agric For Meteorol* 140: 79-96.
- Migliavacca M, Meroni M, Busetto L, Colombo R, Zenone T, Matteucci G, Manca G, Seufert G (2009) Modeling Gross Primary Production of Agro-Forestry Ecosystems by Assimilation of Satellite-Derived Information in a Process-Based Model. *Sensors* 9: 922–942.
- Montagnani L, Manca G, Canepa E, Georgieva E, Acosta M, Feigenwinter C, Janous D, Kerschbaumer G, Lindroth A, Minach L, Minerbi S, Mölder M, Pavelka M, Seufert G, Zeri M, Ziegler W (2009) A new mass conservation approach to the study of CO<sub>2</sub> advection in an alpine forest. *J Geophys Res* 114: D07306.
- Moors E J (2012) Water Use of Forests in The Netherlands. PhD-thesis, Vrije Universiteit Amsterdam, the Netherlands.
- Pilegaard K, Ibrom A, Courtney MS, Hummelshøj P, Jensen NO (2011) Increasing net CO<sub>2</sub> uptake by a Danish beech forest during the period from 1996 to 2009. *Agric For Meteorol* 151: 934–946.
- Rambal S, Joffre R, Ourcival JM, Cavender-Bares J, Rocheteau A (2004) The growth respiration component in eddy CO<sub>2</sub> flux from a *Quercus ilex* Mediterranean forest. *Glob Chang Biol* 10: 1460–1469.
- Rey A, Pegoraro E, Tedeschi V, De Parri I, Jarvis PG, Valentini R (2002) Annual variation in soil respiration and its components in a coppice oak forest in Central Italy. *Glob Chang Biol* 8: 851–866.
- Ruehr NK, Martin JG, Law BE (2012) Effects Of Water Availability On Carbon And Water Exchange In A Young Ponderosa Pine Forest: Above- And Belowground Responses. *Agric For Meteorol* 164: 136-148.
- Sabbatini S, Arriga N, Bertolini T, Castaldi S, Chiti T, Consalvo C, Njakou Djomo S, Gioli B, Matteucci G, Papale D (2016) Greenhouse gas balance of cropland conversion to bioenergy poplar short-rotation coppice. *Biogeosciences* 13: 95–113.
- Saleska SR, Miller SD, Matross DM, Goulden ML, Wofsy SC, Rocha HR da, Camargo PB de, Crill P, Daube BC, Freitas HC de, Hutyrá L, Keller M, Kirchhoff V, Menton M, Munger JW, Pyle EH, Rice AH, Silva H (2003) Carbon In Amazon Forests: Unexpected Seasonal Fluxes And Disturbance-Induced Losses. *Science* 302: 1554-1557.

- Schmidt MW, Torn MS, Abiven S, Dittmar T, Guggenberger G, Janssens IA, Kleber M, Kögel-Knabner I, Lehmann J, Manning DA, Nannipieri P, Rasse DP, Weiner S, Trumbore SE (2011) Persistence Of Soil Organic Matter As An Ecosystem Property. *Nature* 478: 49-56.
- Sun OJ, Campbell J, Law BE, Wolf V (2004) Dynamics Of Carbon Stocks In Soils And Detritus Across Chronosequences Of Different Forest Types In The Pacific Northwest, USA. *Glob Chang Biol* 10: 1470-1481.
- Suni T, Rinne J, Reissell A, Altimir N, Keronen P, Rannik Ü, Maso MD, Kulmala M, Vesala T (2003) Long-term measurements of surface fluxes above a Scots pine forest in Hyytiälä, southern Finland, 1996–2001. *Boreal Environ Res* 8: 287–301.
- Tedeschi V, Rey A, Manca G, Valentini R, Jarvis PG, Borghetti M (2006) Soil respiration in a Mediterranean oak forest at different developmental stages after coppicing. *Glob Chang Biol* 12: 110–121.
- Thum T, Aalto T, Laurila T, Aurela M, Kolari P, Hari P (2007) Parametrization of two photosynthesis models at the canopy scale in a northern boreal Scots pine forest. *Tellus B* 59: 874–890.
- Wolf S, Eugster W, Potvin C, Turner BL, Buchmann N (2011) Carbon sequestration potential of tropical pasture compared with afforestation in Panama. *Glob Chang Biol* 17: 2763–2780.
- Zeller K, Nikolov N (2000) Quantifying Simultaneous Fluxes Of Ozone, Carbon Dioxide And Water Vapor Above A Subalpine Forest Ecosystem. *Environ Pollut* 107: 1-20.
- Zielis S, Etzold S, Zweifel R, Eugster W, Haeni M, Buchmann N (2014) NEP of a Swiss subalpine forest is significantly driven not only by current but also by previous year's weather. *Biogeosciences* 11: 1627–1635.